

Combinatorial Mesh Calculus (CMC): Lecture 12

.....

Lectured by: **Dr. Kiprian Berbatov**¹

Lecture Notes Compiled by: **Muhammad Azeem**¹

Under the supervision of: **Prof. Andrey P. Jivkov**¹

¹ Department of Mechanical and Aerospace Engineering, The University of Manchester, Oxford Road,
Manchester M13 9PL, UK



Area of a Disk via Green's Theorem

Setup

For $r > 0$, let

$$M = \overline{B_r}(0, 0) = \{(x, y) \in \mathbb{R}^2 \mid x^2 + y^2 \leq r^2\},$$
$$\partial M = S_r^1 = \{(x, y) \in \mathbb{R}^2 \mid x^2 + y^2 = r^2\}.$$

The area form on $\mathbb{R}^2$ is $dx \wedge dy$. Since $dx \wedge dy = d(x dy)$,


$$\text{Area}(M) = \int_M dx \wedge dy = \int_M d(x dy) = \int_{\partial M} x dy \quad (\text{Stokes/Green}).$$

Parametrization and Computation

Parametrize ∂M by $\gamma(\phi) = (r \cos \phi, r \sin \phi)$, $\phi \in [0, 2\pi]$. Then $x(\phi) = r \cos \phi$, $y(\phi) = r \sin \phi$, so $dy = r \cos \phi d\phi$ and

$$\begin{aligned}\int_{\partial M} x dy &= \int_0^{2\pi} (r \cos \phi) (r \cos \phi d\phi) \\ &= r^2 \int_0^{2\pi} \cos^2 \phi d\phi = r^2 \cdot \pi \\ &= \pi r^2.\end{aligned}$$

Visualization



Metric Tensor (Riemannian Metric)

Definition (Metric)

Let M be a smooth manifold. A **metric** on M is a bilinear map

$$g : \mathcal{X}(M) \times \mathcal{X}(M) \longrightarrow \mathcal{F}(M),$$

satisfying, for all $X, Y \in \mathcal{X}(M)$:

- *Symmetry*: $g(X, Y) = g(Y, X)$.
- *Positive-definiteness*: $g(X, X) \geq 0$ in $\mathcal{F}(M)$, and $g(X, X) = 0$ iff $X = 0$.

Riemannian manifold

Let M be a smooth manifold and $g : \mathcal{X}(M) \times \mathcal{X}(M) \longrightarrow \mathcal{F}(M)$ be a metric on M . Then the pair (M, g) is a **Riemannian manifold**.

Metric Examples

Examples in Coordinates

Fix local coordinates $(x^1, \dots, x^D)$ and frame $\left\{ \frac{\partial}{\partial x^i} \right\}$.

$$g\left(\frac{\partial}{\partial x^i}, \frac{\partial}{\partial x^j}\right) = g_{ij} \in \mathcal{F}(M), \quad g = \sum_{i,j=1}^D g_{ij} dx^i \otimes dx^j,$$

with the matrix field $\underline{g} = (g_{ij})_{1 \leq i,j \leq D}$ pointwise symmetric positive-definite.

- **Euclidean metric on $\mathbb{R}^D$:** $g_{ij} = \delta_{ij}$, i.e.

$$\underline{g} = I_D = \begin{pmatrix} 1 & 0 & \cdots & 0 \\ 0 & 1 & \cdots & 0 \\ \vdots & \vdots & \ddots & \vdots \\ 0 & 0 & \cdots & 1 \end{pmatrix}, \quad g = \sum_{i=1}^D dx^i \otimes dx^i.$$

Metric Examples

General metric on $\mathbb{R}^D$:

- $\underline{g}(x) = (g_{ij}(x))$ with $g_{ij} = g_{ji}$ and $\underline{g}(x)$ positive-definite for each x .

Historical note:

Bernhard Riemann (1826–1866) introduced Riemannian metrics, founding the intrinsic geometry of manifolds and influencing modern differential geometry and physics.

Flat, Sharp and Dual Metric

Given a Riemannian metric g , the **flat** operator

$$\flat : \mathcal{X}(M) \rightarrow \Omega^1(M), \quad (\flat X)(Y) = g(X, Y),$$

is a $C^\infty(M)$ –isomorphism. Its inverse is the **sharp** operator

$$\sharp = \flat^{-1} : \Omega^1(M) \rightarrow \mathcal{X}(M).$$

They induce a metric on 1–forms,

$$g^* : \Omega^1(M) \times \Omega^1(M) \rightarrow \mathcal{F}(M), \quad g^*(\omega, \eta) = \omega(\sharp \eta).$$

Musical Isomorphisms

Flat/Sharp and Dual Metric

In coordinates, with $g = \sum g_{ij} dx^i \otimes dx^j$ and inverse matrix (g^{ij}) :

$$g^* = \sum_{i,j=1}^D g^{ij} \frac{\partial}{\partial x^i} \otimes \frac{\partial}{\partial x^j}.$$

$$\begin{array}{ccc} \mathcal{X}(M) & \xleftarrow{\flat} & \Omega^1(M) \\ & \sharp = \flat^{-1} & \\ g(X, Y) = (\flat X)(Y), & g^*(\omega, \eta) = \omega(\sharp \eta). & \end{array}$$

Pullback Metric on a Submanifold

Definition (Induced Metric)

Let (M, g) be Riemannian and $S \subseteq M$ a smooth submanifold with embedding $\iota : S \hookrightarrow M$, $\iota(s) = s$. The **induced (pullback) metric** on S is

$$\iota^*g = \sum_{i,j=1}^D (g_{ij} \circ \iota) d(x^i \circ \iota) \otimes d(x^j \circ \iota).$$

Interpretation

Express g in the ambient coordinates, then pull back each coefficient and each coordinate differential to S via ι (i.e. *change of variables* to the coordinates on S).

Example

Unit Sphere $S^2 \subset \mathbb{R}^3$

Let $M = B_1(0, 0, 0) = \{(x, y, z) \mid x^2 + y^2 + z^2 \leq 1\}$ and $B_1(0, 0, 0) \subset \mathbb{R}^3$, $S = \partial M = S^2 = \{(x, y, z) \mid x^2 + y^2 + z^2 = 1\}$ and $g = dx \otimes dx + dy \otimes dy + dz \otimes dz$ (Euclidean).

In spherical coordinates on S^2 ,

$$x = \sin \theta \cos \phi, \quad y = \sin \theta \sin \phi, \quad z = \cos \theta,$$

with

$$(\theta, \phi) \in (0, \pi) \times (0, 2\pi).$$

Compute

$$dx = \cos \theta \cos \phi d\theta - \sin \theta \sin \phi d\phi,$$

$$dy = \cos \theta \sin \phi d\theta + \sin \theta \cos \phi d\phi,$$

$$dz = -\sin \theta d\theta.$$

Example

Unit Sphere $S^2 \subset \mathbb{R}^3$

Then

$$\iota^*g = dx^2 + dy^2 + dz^2 = (d\theta)^2 + \sin^2 \theta (d\phi)^2.$$


Hence, in the chart (θ, ϕ) ,

$$\underline{h} = \begin{pmatrix} 1 & 0 \\ 0 & \sin^2 \theta \end{pmatrix}, \quad \underline{h}^{-1} = \begin{pmatrix} 1 & 0 \\ 0 & \frac{1}{\sin^2 \theta} \end{pmatrix},$$

and

$$h^{-1} = \frac{\partial}{\partial \theta} \otimes \frac{\partial}{\partial \theta} + \frac{1}{\sin^2 \theta} \frac{\partial}{\partial \phi} \otimes \frac{\partial}{\partial \phi}.$$

Visualization



$$\iota^*g = d\theta^2 + \sin^2 \theta d\phi^2 \text{ on } S^2$$

Orthonormal Frames

Definition

Let (M, g) be a D -dimensional Riemannian manifold. A frame $(X_1, \dots, X_D)$ in $\mathcal{X}(M)$ is **orthonormal** if

$$g(X_i, X_j) = \delta_{ij} \quad (1 \leq i, j \leq D).$$

The dual coframe (or dual basis) $(\omega^1, \dots, \omega^D)$ in $\Omega^1(M)$ is orthonormal for g^* .

Examples of Orthonormal Bases

Example 1: Euclidean Space $(\mathbb{R}^D, g = \sum_{i=1}^D dx^i \otimes dx^i)$

Consider the standard Cartesian coordinates $(x^1, \dots, x^D)$ on $\mathbb{R}^D$. The standard basis of vector fields is

$$\left(\frac{\partial}{\partial x^1}, \frac{\partial}{\partial x^2}, \dots, \frac{\partial}{\partial x^D} \right) \in \mathcal{X}(\mathbb{R}^D),$$

and the corresponding dual basis of 1-forms is

$$(dx^1, dx^2, \dots, dx^D) \in \Omega^1(\mathbb{R}^D).$$

For this metric,

$$g\left(\frac{\partial}{\partial x^i}, \frac{\partial}{\partial x^j}\right) = \delta_{ij}, \quad g^*(dx^i, dx^j) = \delta_{ij}.$$

Examples of Orthonormal Bases

Example 1: Euclidean Space ($\mathbb{R}^D, g = \sum_{i=1}^D dx^i \otimes dx^i$)

Hence both sets are orthonormal:


$$g\left(\frac{\partial}{\partial x^i}, \frac{\partial}{\partial x^i}\right) = 1, \quad g\left(\frac{\partial}{\partial x^i}, \frac{\partial}{\partial x^j}\right) = 0 \quad (i \neq j),$$

and similarly for the dual coframe.

Interpretation:

The Euclidean metric measures inner products as the usual dot product; hence, in these coordinates, vectors and covectors have the same orthonormal structure.

Visualization



Orthonormal frame in $\mathbb{R}^3$

Example 2: Orthonormal Basis

On the Sphere S^2

On S^2 with the induced metric from $\mathbb{R}^3$,

$$h = d\theta^2 + \sin^2 \theta d\phi^2,$$

where (θ, ϕ) are the standard spherical coordinates:

$$x = \sin \theta \cos \phi, \quad y = \sin \theta \sin \phi, \quad z = \cos \theta.$$

Compute h on the coordinate vector fields:

$$h\left(\frac{\partial}{\partial \theta}, \frac{\partial}{\partial \theta}\right) = 1,$$

$$h\left(\frac{\partial}{\partial \phi}, \frac{\partial}{\partial \phi}\right) = \sin^2 \theta,$$

Example 2: Orthonormal Basis

On the Sphere S^2

$$h\left(\frac{\partial}{\partial\theta}, \frac{\partial}{\partial\phi}\right) = 0.$$

Hence, the rescaled frame


$$\left(\frac{\partial}{\partial\theta}, \frac{1}{\sin\theta} \frac{\partial}{\partial\phi}\right)$$

is orthonormal:

$$h\left(\frac{\partial}{\partial\theta}, \frac{\partial}{\partial\theta}\right) = h\left(\frac{1}{\sin\theta} \frac{\partial}{\partial\phi}, \frac{1}{\sin\theta} \frac{\partial}{\partial\phi}\right) = 1, \quad h\left(\frac{\partial}{\partial\theta}, \frac{1}{\sin\theta} \frac{\partial}{\partial\phi}\right) = 0.$$

The dual orthonormal coframe is $(d\theta, \sin\theta d\phi)$.

Example 2: Orthonormal Basis



Orthonormal tangent frame and coframe on S^2

Inner Product on Differential Forms

Definition

Let (M, g) be a D -dimensional Riemannian manifold. The metric g induces an inner product on p -forms:

$$g_p^* : \Omega^p(M) \times \Omega^p(M) \rightarrow \mathcal{F}(M), \quad p = 0, 1, \dots, D,$$

defined recursively:

$$g_0^*(f_1, f_2) = f_1 f_2, \quad f_1, f_2 \in \mathcal{F}(M),$$

$$g_1^*(\omega, \eta) = g^*(\omega, \eta), \quad \omega, \eta \in \Omega^1(M),$$

Inner Product on p -Forms

Definition

Let (M, g) be a D -dimensional Riemannian manifold. For $p \geq 2$, the metric g induces an inner product on p -forms

$$g_p^* : \Omega^p(M) \times \Omega^p(M) \longrightarrow \mathcal{F}(M),$$

defined on decomposable forms as

$$g_p^*(\omega_1 \wedge \cdots \wedge \omega_p, \eta_1 \wedge \cdots \wedge \eta_p) = \begin{vmatrix} g^*(\omega_1, \eta_1) & g^*(\omega_1, \eta_2) & \cdots & g^*(\omega_1, \eta_p) \\ g^*(\omega_2, \eta_1) & g^*(\omega_2, \eta_2) & \cdots & g^*(\omega_2, \eta_p) \\ \vdots & \vdots & \ddots & \vdots \\ g^*(\omega_p, \eta_1) & g^*(\omega_p, \eta_2) & \cdots & g^*(\omega_p, \eta_p) \end{vmatrix}.$$

The determinant encodes all mutual inner products between $\{\omega_i\}$ and $\{\eta_j\}$:

$$g_p^*(\omega_1 \wedge \dots \wedge \omega_p, \eta_1 \wedge \dots \wedge \eta_p) = \sum_{\sigma \in S_p} \text{sgn}(\sigma) \prod_{k=1}^p g^*(\omega_k, \eta_{\sigma(k)}),$$

where S_p is the permutation group on $\{1, \dots, p\}$ and $\text{sgn}(\sigma)$ is the sign of permutation σ . This generalizes the usual Euclidean determinant form of the inner product on wedge products.

$$g_p^*(\omega_1 \wedge \dots \wedge \omega_p, \eta_1 \wedge \dots \wedge \eta_p) = \det[g^*(\omega_i, \eta_j)]_{1 \leq i, j \leq p}$$


$$g_p^*(\omega, \eta) = \sum_{\sigma \in S_p} \text{sgn}(\sigma) g^*(\omega_1, \eta_{\sigma(1)}) \cdots g^*(\omega_p, \eta_{\sigma(p)}).$$

Example

S^2 with $h = d\theta^2 + \sin^2 \theta d\phi^2$

$$\begin{aligned} g_2^*(d\theta \wedge d\phi, d\theta \wedge d\phi) &= \det \begin{pmatrix} g^*(d\theta, d\theta) & g^*(d\theta, d\phi) \\ g^*(d\phi, d\theta) & g^*(d\phi, d\phi) \end{pmatrix} \\ &= \det \begin{pmatrix} 1 & 0 \\ 0 & \frac{1}{\sin^2 \theta} \end{pmatrix} = \frac{1}{\sin^2 \theta}. \end{aligned}$$

Hence $g_2^*(\sin \theta d\theta \wedge d\phi, \sin \theta d\theta \wedge d\phi) = \sin^2 \theta \cdot \frac{1}{\sin^2 \theta} = 1.$



$$g_2^*(d\theta \wedge d\phi, d\theta \wedge d\phi) = \frac{1}{\sin^2 \theta}$$

Volume Form

Definition

Let (M, g) be a D -dimensional Riemannian manifold. Since $\Omega^D(M)$ is 1-dimensional, there exist (up to sign) two D -forms ω and $-\omega$ such that

$$g_D^*(\omega, \omega) = 1.$$

Any such form is a **volume form** of (M, g) . If M is oriented, the sign of ω is fixed canonically, and we denote $\text{vol}_g = \omega$.

Examples

Example

- On $\mathbb{R}^D$ with $g = \sum dx^i \otimes dx^i$:

$$\text{vol}_g = \pm dx^1 \wedge dx^2 \wedge \cdots \wedge dx^D.$$

- On S^2 with $h = d\theta^2 + \sin^2 \theta d\phi^2$:

$$\text{vol}_h = \sin \theta d\theta \wedge d\phi.$$

Proposition

Statement: Volume Form from Orthonormal Basis

Let (M, g) be an oriented D -dimensional Riemannian manifold and $e^1, \dots, e^D$ an orthonormal basis of $\Omega^1(M)$ consistent with the orientation. Then

$$\text{vol}_g = e^1 \wedge e^2 \wedge \cdots \wedge e^D.$$

Examples


Example

- On $(\mathbb{R}^D, g)$ with standard basis $dx^1, \dots, dx^D$:

$$\text{vol}_g = dx^1 \wedge dx^2 \wedge \dots \wedge dx^D.$$

- On S^2 with orthonormal coframe $(d\theta, \sin \theta d\phi)$:

$$\text{vol}_h = d\theta \wedge (\sin \theta d\phi) = \sin \theta d\theta \wedge d\phi.$$



$$\text{vol} = e^1 \wedge e^2$$

Hodge Star Operator

Definition

Let (M, g) be a D -dimensional oriented Riemannian manifold with volume form vol_g . The **Hodge star** is the unique $\mathcal{F}(M)$ -linear map

$$\star_p : \Omega^p(M) \rightarrow \Omega^{D-p}(M)$$

such that for all $\omega, \eta \in \Omega^p(M)$,

$$g_p^*(\omega, \eta) \text{vol}_g = \omega \wedge \star_p \eta.$$

Equivalently, for $\omega \in \Omega^p(M)$, $\eta \in \Omega^{D-p}(M)$,

$$g_{D-p}^*(\star_p \omega, \eta) \text{vol}_g = \omega \wedge \eta.$$

Theorem: Properties of the Hodge Star

Statement

Let (M, g) be an oriented D -dimensional Riemannian manifold and $e^1, \dots, e^D$ be an oriented orthonormal basis of $\Omega^1(M)$.

1. For an index set $I = \{i_1 < \dots < i_p\}$, let $e^I = e^{i_1} \wedge \dots \wedge e^{i_p}$, and J the complementary set of indices. Then

$$\star_p e^I = s e^J, \quad s \in \{-1, 1\},$$

where $e^I \wedge e^J = s \operatorname{vol}_g = s e^1 \wedge \dots \wedge e^D$.

2. $\star_{D-p} \circ \star_p = (-1)^{p(D-p)} \operatorname{id}_{\Omega^p(M)}$.
In particular, if D is odd, $\star_{D-p} \circ \star_p = \operatorname{id}_{\Omega^p(M)}$.


Examples: Hodge Star in $D = 2$

Example: S^2 with orthonormal coframe
($e^1 = d\theta$, $e^2 = \sin \theta d\phi$)

$$\star_0 (1) = e^1 \wedge e^2 = \text{vol},$$

$$\star_1 (e^1) = e^2, \quad \star_1 (e^2) = -e^1,$$

$$\star_2 (\text{vol}) = \star_2 (e^1 \wedge e^2) = 1.$$



$$\star_1 e^1 = e^2, \quad \star_1 e^2 = -e^1$$

Examples: Hodge Star in $D = 3$

Example: $\mathbb{R}^3$ with orthonormal coframe


$$(e^1 = dx, e^2 = dy, e^3 = dz)$$

$$\star_0(1) = e^1 \wedge e^2 \wedge e^3 = \text{vol},$$

$$\star_1(e^1) = e^2 \wedge e^3, \quad \star_1(e^2) = e^3 \wedge e^1, \quad \star_1(e^3) = e^1 \wedge e^2,$$

$$\star_2(e^1 \wedge e^2) = e^3, \quad \star_2(e^2 \wedge e^3) = e^1, \quad \star_2(e^3 \wedge e^1) = e^2,$$

$$\star_3(e^1 \wedge e^2 \wedge e^3) = 1.$$



Summary

- Area of a disk via Green/Stokes:

$$\text{Area}(\overline{B_r}) = \int_{\partial B_r} x \, dy = \pi r^2.$$

- Riemannian metric g gives lengths/angles:

$$g = \sum g_{ij} \, dx^i \otimes dx^j, \text{ with inverse } g^* = \sum g^{ij} \, \partial_i \otimes \partial_j.$$

- Musical isomorphisms $\flat, \sharp$ identify vectors and 1-forms.
- Induced metric on submanifolds: $\iota^* g$; for S^2 , $d\theta^2 + \sin^2 \theta \, d\phi^2$.
- Orthonormal frames simplify computations (e.g. on S^2 : $(\partial_\theta, \frac{1}{\sin \theta} \partial_\phi)$).
- g_p^* defines inner products on all p -forms via determinant of $g^*(\cdot, \cdot)$.

Summary

- Volume form vol_g satisfies $g_D^*(\text{vol}_g, \text{vol}_g) = 1$ and encodes orientation.
- $\text{vol}_g = e^1 \wedge \cdots \wedge e^D$ for any oriented orthonormal coframe.
- Hodge star $\star_p$ maps $\Omega^p(M) \rightarrow \Omega^{D-p}(M)$ satisfying $g_p^*(\omega, \eta)\text{vol}_g = \omega \wedge \star_p \eta$.
- Key identity: $\star_{D-p} \circ \star_p = (-1)^{p(D-p)} \text{id}$.

Thanks

